

Treatment of Land-Use and Urbanization

LECTURE 9



Alexander Mahura

NetFAM School and Workshop on
“Integrated Modelling of Meteorological and
Chemical Transport Processes / Impact of
Chemical Weather on Numerical Weather
Prediction and Climate Modelling”

Saint-Petersburg, Russia
7-15 July 2008



 **norden**
NetFAM/NordForsk
NetFAM:

Nordic Network on Fine-scale Atmospheric Modelling

*Danish Meteorological Institute, DMI,
Copenhagen, Denmark*

NetFAM Summer School on Integrated Modelling ...
7-12 July 2008, Zelenogorsk, Russia

Outline of the Lecture

- Land-cover and land-use: classification, datasets, etc.;

-
- Urban lands: some statistics;

- Urbanized areas: urban boundary layer, features, controls, characteristics, approaches for treatment, etc.

- Urbanization of Enviro-HIRLAM: modules, urban districts, anthropogenic heat flux, some results (on example of Copenhagen Metropolitan area, Denmark), applicability, etc.

An aerial photograph of a rural landscape. A river flows through the center, surrounded by green fields. In the background, a town is visible, and there are patches of yellow fields. The text "LAND COVER and LAND USE" is overlaid in red with a white outline. There are decorative teal and light blue shapes on the left side of the image.

LAND COVER and LAND USE



Land Cover and Land Use

Land cover -

defined as observed physical cover, as seen from the ground or through remote sensing, including natural or planted vegetation and human constructions (buildings, roads, etc.) which cover the earth's surface. Water, ice, bare rock or sand surfaces count as land cover.

Land Use –

defined as a series of activities undertaken to produce one or more goods or services. A given land use may take place on one or several pieces of land, and several land uses may occur on the same piece of land.



Why There is a Need for Meteorological Modelling

Simulate exchanges between surface and atmosphere (momentum, heat, water, chemical species, etc.);


Take into account the climate variability from one region to another

Separate the surface schemes from the atmospheric model - allows to use the same surface code for several atmospheric models (NWP models runs)
- easy switch between surface schemes and options;

All surface fields necessary to land surface schemes


CORINE: Coordination of Information on the Environment

<http://www.eea.eu.int>



**LCDB - based on interpretation of satellite images for 1989 and 1990,
land cover types in 44 standard classes,
GIS ARC/INFO format, at an original scale of 1:100,000 (consistent and comparable
with similar land cover databases in other European countries.
Update - 2000**

CORINE : EU Countries




Land cover map - CORINE dataset

250 m resolution, 44 classes

- | | |
|--|--|
|  Continuous urban fabric |  Agro-forestry areas |
|  Discontinuous urban fabric |  Broad-leaved forest |
|  Industrial or commercial units |  Coniferous forest |
|  Road and rail networks and associated land |  Mixed forest |
|  Port areas |  Natural grasslands |
|  Airports |  Moors and heathland |
|  Mineral extraction sites |  Sclerophyllous vegetation |
|  Dump sites |  Transitional woodland-shrub |
|  Construction sites |  Beaches, dunes, sands |
|  Green urban areas |  Bare rocks |
|  Sport and leisure facilities |  Sparsely vegetated areas |
|  Non-irrigated arable land |  Burnt areas |
|  Permanently irrigated land |  Glaciers and perpetual snow |
|  Rice fields |  Inland marshes |
|  Vineyards |  Peat bogs |
|  Fruit trees and berry plantations |  Salt marshes |
|  Olive groves |  Salines |
|  Pastures |  Intertidal flats |
|  Annual crops associated with permanent crops |  Water courses |
|  Complex cultivation patterns |  Water bodies |
|  Land principally occupied by agriculture, with significant areas of natural vegetation |  Coastal lagoons |
| |  Estuaries |
| |  Sea and ocean |
| |  NODATA |

CORINE : Denmark : Classification



**Denmark: Land cover map - CORINE dataset
(21 class, Sattler, 1999)**

Description of land-class

Crops, Mixed Farming
Irrigated Crops
Bogs and Marshes
Evergreen Needle-leaf Trees
Deciduous Needle-leaf Tree
Deciduous Broad-leaf Trees
Evergreen Broad-leaf Trees
Evergreen Shrubs
Deciduous Shrubs
Interrupted Forest
Mixed Forest
Tundra
Short Grass
Tall Grass
Desert
Semi-desert
Ocean
Inland Water
Water and Land Mixtures
Ice Caps and Glaciers

Urban area

ECOCLIMAP, USGS, PELCOM, etc.



Sea	Port and leisure facilities
Continuous urban fabric	Non-irrigated arable land
Discontinuous urban fabric	Permanently irrigated land
Industrial and commercial units	Rice fields
Road and rail networks and assoc	Vineyards
Port areas	Fruit trees and berry plantation
Airports	Olive groves
Mineral extraction sites	Pastures
Dump sites	Annual crops associated with perman
Construction sites	Complex cultivation patterns
Green urban areas	Land principally occupied by agr
Agro-forestry areas	Burnt areas
Broad-leaved forest	Glaciers and perpetual snow
Coniferous forest	Inland marshes
Mixed forest	Peat bogs
Natural grasslands	Salt marshes
Moors and heath lands	Salines
Sclerophyllous vegetation	Intertidal flats
Transitional woodland-scrub	Water courses
Beaches, sand, dunes	Water bodies
Bare rocks	Coastal lagoons
Sparsely vegetated areas	Estuaries

**Land cover map - ECOCLIMAP dataset
1 km resolution**


+ Other datasets, USGS, PELCOM, etc.

URBAN LANDS



EU: Urban Land Development


EU statistics from <http://www.eea.eu.int>



EU: Urban Land Uptake by Origin


During 1990-2000 from all areas converted to artificial land-use

Origin of urban land uptake as percentage of total uptake




EU: Urban Land Uptake by Countries

Urban land uptake by countries



EU: Urban Land Uptake by Metropolitan Areas



**Urban land
uptake
by megacities**

Urban and infrastructure development

- 1 to 5 %
- 5 to 10 %
- More than 10 %

Urban zones 1990

- More than 50 000 people

Green background index

- 0 to 60 %
- 61 to 100 %

GLOBAL: climate change

URBANIZED AREAS

REGIONAL:
acid rain,
tropospheric ozone,
aerosols, greenhouse gases

LOCAL:
air pollution
health effects


Heat island

plume

people

Urban Boundary Layer

Urban Boundary Layer, UBL vs. 'rural' homogeneous boundary layer is characterised by greatly enhanced mixing, resulting from both the large surface roughness and increased surface heating, and by horizontal inhomogeneity of meteorological fields due to variations in surface roughness and heating from rural, sub-urban, to central areas of cities.





Features for Urban Areas

- **Local-scale inhomogeneties, sharp changes of roughness and heat fluxes;**
- **Wind velocity reduce effect due to buildings;**
- **Redistribution of eddies due to buildings, from large to small;**
- **Trapping of radiation in street canyons;**
- **Effect of urban soil structure, diffusivities heat and water vapour;**
- **Anthropogenic heat fluxes, urban heat island;**
- **Internal urban boundary layers, urban mixing height,**
- **Effects of pollutants (aerosols) on urban meteorology and climate;**
- **Urban effects on clouds, precipitation and thunderstorms.**

Controls on Urban Climate Effects (including Urban Heat Island)

(Oke et al., 1980)

Fixed

City Location

- climate
- topography
- rural surrounds

City Size

- fetch distance
- density of use

Time

- day
- season

Modulators

Weather

- wind
- cloud
- stability

City Form


- fabric
- structure
- cover

City Metabolism

- energy/water use
- emissions

Controllable

(policy, planning, design)





Surface Energy Balance, Characteristics of Urban Surfaces

$$Q^* = K\downarrow - K\uparrow + L\downarrow - L\uparrow = Q_H + Q_E + \Delta Q_S + Q \text{ [W/m}^2\text{]}$$

Q^* - net all-wave radiation;
 $K\downarrow$ and $K\uparrow$ - incoming and outgoing reflected shortwave radiation;
 $L\downarrow$ and $L\uparrow$ - incoming and outgoing longwave radiation;
 Q_H and Q_E - turbulent sensible and latent heat fluxes,
 ΔQ_S - storage heat flux,
 Q - other sources/sinks.

- **Altered albedo – can be higher or lower,**
- **Higher heat capacity,**
- **Lower moisture flux to atmosphere,**
- **Larger roughness elements,**
- **Increased surface area,**
- **Source of anthropogenic heat and emissions,**
- **Impermeable to water,**
- **Decreased net longwave radiation loss.**



Approaches for Treatment of Urban Boundary Layer Features


- **Urban roughness effects**
(*Bornstein, 1975, 2001; Hunt et al., 2003*)
- **Urban surface energy balance**
(*Oke et al., 1999; Piringer et al., 2002*)
- **Town Energy Balance scheme**
(*Masson, 2000*)
- **Urban surface exchange sub-layer model**
(*Martilli et al., 2002*)
- **Soil model for sub-meso scales urban version**
(*Dupont et al., 2006ab*)
- **Prognostic equations for UBL height**
(*Zilitinkevich et al., 2002+; Gryning and Bartchvarova, 2002*).

URBANIZATION of Enviro-HIRLAM





Enviro-HIRLAM: Urbanization Modules

- anthropogenic heat flux and roughness (**AHF+R**)
- building effect parameterization (**BEP**)
- soil model for sub-meso scales urban version (**SM2-U**)



Enviro-HIRLAM : Urban : Land Surface Scheme, Tiles and Urban Areas, Modelling Domains, and Focus

- Land surface scheme: Interaction Soil-Biosphere-Atmosphere (ISBA)
- Tiles (low vegetation, forest, ice, snow, water, bare soil) + urban fraction
- High resolution domains: -U01/-I01 (horiz resol of 1.4 km)
- Climate Generation Files, + surface and meteorology related data
- Focus: Copenhagen metropolitan area (Island of Sjealland)



Urban Districts : Classification

Residential (RD)



City Center/High Buildings District (CC/HBD)



GIS

Industrial Commercial (ICD)



Extraction of districts related characteristics (statistics):

Morphology parameters (avg. height, volume, perimeter, compactness, space between buildings),

Cover modes (surface density (SD) of buildings, of vegetation, hydrography, roads, N buildings),

Aerodynamic parameters (roughness length, displacement height, frontal and lateral SD),

Anthropogenic Heat Flux in Urban Areas

can be calculated based on assumption of dependency/ proportionality to other urban characteristics:


1. Population density maps with a high resolution in urban areas;
2. Satellite images of the night lightness over urban areas (but difficulties to use for industrial and developing countries, should be corrected);
3. Land-use classification as a percentage of urban classes (central part, urban, sub-urban, industrial, etc.);
4. Emission inventory for specific pollutants typical for urban areas (e.g., due to traffic emission, etc.);
5. Monitoring or simulation of concentration fields for specific air pollutants typical for urban areas.



Reference avg. value: up to 100 W/m²


BEP : Building Effect Parameterization

(Martilli et al., 2002)




MeteoModelling: BEP Module

Difference between runs: 01 Aug 2004, 06 UTC



(control vs. urbanized run)
Difference field for temperature at 2 m


(control vs. urbanized run)
Difference field for wind at 10 m




MeteoModelling: BEP Module



Urban station N-6180: Copenhagen area : Verification



July 2004




August 2004

Diurnal variability for 00 UTC forecasts for the average wind velocity at 10 m for the urban station N-6180 in the Copenhagen metropolitan area as function of the forecast length based on the DMI-HIRLAM-I01+**BEP /U0D/** and -I01-CTRL /C0D/ model runs vs. **observations**

SM2-U : Soil Model for Sub-Meso scales

Urbanized version : Thermal budget


(Dupont et al., 2006ab)



SM2-U : Soil Model for Sub-Meso scales

Urbanized version : Water budget





(Dupont et al., 2006ab)



Revised Land Use Classification : SM2-U Module






BARE	Bare soil without vegetation
NAT	Bare soil located between sparse vegetation elements
VEGN	Vegetation over bare soil
VEGA	Vegetation over paved surfaces
ART	Paved surfaces located between the sparse vegetation elements
BAT	Building/roofs
EAU	Water surfaces



Dominated type

MeteoModelling: SM2-U Module



Metropolitan area : Copenhagen : Surface temperature



MeteoModelling: SM2-U Module



Metropolitan area : Copenhagen : Surface temperature : Urban districts




CC/HBD	City center / high buildings district
ICD	Industrial commercial district
RD	Residential district
Non-urban	Non-urban areas (no BAT type)

MeteoModelling: SM2-U Module



Metropolitan area : Copenhagen : Storage and sensible heat fluxes : Types of surfaces



BARE	Bare soil without vegetation
NAT	Bare soil located between sparse vegetation elements
VEGN	Vegetation over bare soil
VEGA	Vegetation over paved surfaces
ART	Paved surfaces located between the sparse vegetation elements
BAT	Building/roofs
EAU	Water surfaces



Evaluation of Results : Items

- Specific dates/ short- and long-term periods selected,

- Diurnal cycle,
- Month-to-month variability,
- Difference between the control vs. urban runs,
- Meteorological variables of key importance,
- Urban districts of different nature:
 - City Center,
 - High Buildings District,
 - Industrial Commercial District,
 - Residential District,
- Types of surfaces (including urban variants),
- Focus: impact of urban areas on simulated meteorological fields

Urbanization: Applicability of Results

Testing and verification of numerical weather prediction and climatological models performance over high resolution model domains, and especially, over the urbanized areas;

- Investigation of temporal and spatial variability of various meteorological and derived variables over urbanized areas;
- Improvements in land use classification and climate generation properties;
- Distinguishing and selection of types of urban districts and their properties;
- Urbanization of climate regional and global models.



Thanks

- DMI FM HIRLAM and Environmental Meteorology Groups, the ECN (France) and EPFL (Switzerland) research teams,
- FUMAPEX, COST Actions, MEGAPOLI, etc.